

**Research Article****Potential for single cell protein production by yeast isolated from fermented foods**

Viet Hoang Quoc<sup>1,2</sup>, Dung Le Thi Hoang<sup>1,2</sup>, Loc Pham Thanh<sup>1,2</sup>, Hue Nguyen Thi<sup>1,2</sup>, Hue Pham Thi<sup>1</sup>,  
Minh Nguyen Hong<sup>1,2</sup>, Phong Tran Huu<sup>1,2\*</sup>

<sup>1</sup>Bioresources Research Center, Phenikaa School of Engineering,  
Phenikaa University, Hanoi, Vietnam

<sup>2</sup>Faculty of Biotechnology, Chemistry and Environmental Engineering, Phenikaa School of Engineering,  
Phenikaa University, Hanoi, Vietnam

(Received: 30 Nov 2025; Revised: 12 Dec 2025; Accepted: 31 Mar 2026)

**Abstract**

In this study, 36 yeast strains were isolated from fermented foods collected in different areas of Northern Vietnam. The screening process yielded 15/36 strains capable of good growth with Cell Dry Weight (CDW) ranging from 10-14 g/L, and 10/36 strains having the potential for single cell protein production with Total Protein Content (TPC) exceeding 57% (g/100 g cell dry weight) after 24 h of cultivation in YPD liquid medium. The yeast strain YPI-Y2 (with CDW reaching 14.03 g/L; TPC reaching 62.3%) was identified as closely related to the species *Pichia kudriavzevii* (99.77%) based on ITS rDNA sequence analysis and comparison. The nutritional and cultivation characteristics of the *P. kudriavzevii* YPI-Y2 strain, oriented towards cell biomass production and protein accumulation, were determined. The optimal cultivation temperature and initial medium pH condition for the growth and protein accumulation of the YPI-Y2 yeast strain were 30°C and 6.5, respectively, while the suitable carbon source was identified as D-fructose. Simultaneously, this yeast strain maintained good growth and protein accumulation capability on inorganic nitrogen sources (except urea, with 35.62% of TPC). Under suitable cultivation and nutritional conditions, the CDW and TPC values reached 16.12 g/L and 66.62% (g/100 g CDW), respectively.

**Keywords:** Single cell protein (SCP), *Pichia kudriavzevii*, yeast, fermented food, YPI-Y2

**1. INTRODUCTION**

Climate change, coupled with population growth, constitutes a primary determinant of the stability and sustainability of the global food security system. The global population is projected to reach a threshold of approximately 9.3 to 10 billion by 2050. Consequently, addressing the escalating demand for food to sustain human life has become an urgent imperative. Concurrently, nutritional instability poses severe risks to human health, leading to adverse consequences such as malnutrition and muscle atrophy. Collectively, achieving equilibrium between food production and consumption poses a formidable challenge for traditional food production systems that are heavily reliant on agriculture [1]. According to the Food and Agriculture Organization of the United Nations (FAO), human nutritional requirements are categorized into 3 primary macronutrients: fats, carbohydrates, and proteins. Among these, protein is considered the most critical and essential nutrient, given its ubiquitous involvement in virtually all cellular functions and physiological processes, distinguishing it from fats and carbohydrates. Consequently, the exploration of alternative protein sources that are non-animal and non-plant in origin, specifically those that are cost-effective and independent of arable land and freshwater resources, represents a vital strategy in the current global context. In this regard, single-cell protein (SCP) derived from microorganisms emerges as a potent solution. SCP satisfies the rigorous

\*Corresponding author: Tran Huu Phong (E-mail: [phong.tranhuu@phenikaa-uni.edu.vn](mailto:phong.tranhuu@phenikaa-uni.edu.vn))  
<https://doi.org/10.47866/2615-9252/vjfc.4672>

Copyright © 2025 The author(s). This article is licensed under [CC BY-NC 4.0](https://creativecommons.org/licenses/by-nc/4.0/)

criteria established to address food security challenges, being nutrient-dense with minimal reliance on land and freshwater, while simultaneously mitigating environmental impacts [1].

SCP refers to protein-rich microbial biomass produced from various microorganisms, including yeast, filamentous fungi, algae, and bacteria [1]. Among these, yeast-derived single-cell protein (SCP-Y) represents a particularly promising avenue for both research and industrial application [1, 2]. This is primarily because SCP-Y boasts a crude protein content of up to 60% of its dry cell weight, significantly surpassing conventional protein sources such as meat (45%), milk (25%), and soybean (35%) [2]. The issue is worth discussing, and the primary rationale for our selection of yeast as the subject for SCP production is its comprehensive amino acid profile. SCP-Y contains all 8 essential amino acids that the human cannot synthesize *de novo*, including isoleucine (Ile), leucine (Leu), lysine (Lys), phenylalanine (Phe), methionine (Met), threonine (Thr), tryptophan (Trp), and valine (Val). A notable example is lysine. A study by Jach et al. (2022) reported that yeast can synthesize lysine at concentrations up to 70 mg/g protein, which is 2.5 times higher than the 28 mg/g typically found in wheat [2]. These findings underscore the potential of yeast SCP as a viable solution to global nutritional challenges. It offers a low-cost, scalable production method that is independent of arable land and freshwater resources, thereby supporting sustainable food security and minimizing environmental footprints. Consequently, microbial SCP in general, and SCP-Y in particular, is currently garnering significant attention and being intensively exploited worldwide, including in Vietnam [1, 2].

In Vietnam, yeast has been present in many food products for a long time, and has mostly focused on alcoholic beverages. Early research on SCP was carried out by Pham Thanh Ho using oyster mushroom (*Pleurotus*) as a producer, but not yeast [3]. Due to the advantages such as popularity and safety, some researchers have chosen *Saccharomyces cerevisiae* as a SCP potential producer. Trang et al. (2028) optimized biomass production on cost-effective substrates such as molasses, establishing a promising basis for protein-enriched biomass applicable to food and animal feed industries [4]. Hai et al. (2024) utilized resultant seaweed hydrolysate as a nutrient for *S. cerevisiae* biomass production, resulting in  $1.2 \times 10^6$  CFU/mL after 72 h of cultivation [5]. By-product of pulp industry was hydrolysed for SCP production by *Candida utilis* at a pilot scale of 1000 liters for animal feed applications [6]. *Pichia kudriavzevii* is a particularly non-conventional yeast. Although the safety of *P. kudriavzevii* remains controversial, it is found in many fermented foods. Therefore, *P. kudriavzevii* is attracting attention for food and biotechnology applications [7].

In this study, we isolated and screened yeast strains with the capability of rapid growth and concomitant high protein accumulation. Subsequently, one superior strain was selected to investigate its biological characteristics for SCP production applications. The results of this research contribute to the effective exploration and exploitation of non-*Saccharomyces* yeast diversity for SCP production, proposing an effective alternative to animal and plant-based protein in the current context.

## 2. MATERIALS AND METHODS

### 2.1. Materials

A total of five distinct fermented food samples were collected directly from local household producers (Table 1). The samples were subsequently transported to the Bioresource Research Center, Phenikaa University, and stored at 4°C before analysis.

*Table 1. Sample informations for yeast isolation*

Sample name (code)	Sample information (location, collector, ...)
Vegetable pickle (YPI)	Location: Bui Xuong Trach, Hanoi, Vietnam Collector: Le Thi Hoang Dung
“Tré” (fermented meat) (YFPS)	Location: Bac Ninh, Vietnam Collector: Hoang Quoc Viet
Kimchi (YKCS)	Location: Ha Loc, Phu Tho, Vietnam Collector: Hoang Quoc Viet
Fermented grape (NHC-V)	Location: Hoai Duc, Hanoi, Vietnam Collector: Tran Huu Phong
Grape pomace (BN-V)	Location: Ha Thach, Phu Tho, Vietnam Collector: Hoang Quoc Viet

## 2.2. Chemicals and reagents

Standard chemicals: Methyl red (CAS: 493-52-7), and bromocresol green (CAS: 76-60-8, Xilong). Other reagents and solvents consisted of: yeast extract, peptone-r, D-glucose, D-fructose, 98% sulfuric acid, sodium hydroxide, boric acid, potassium sulfate, copper sulfate pentahydrate, and double-distilled water (ddH<sub>2</sub>O).

## 2.3. Equipment and apparatus

General laboratory equipment included: Class II-Biological Safety Cabinet, laminar flow cabinet, 4°C refrigerator, microbial stock storage unit, autoclave, fume-hood, optical microscope, shaking incubator, and cooled incubator. Specialized instrumentation included: Biochrom WPA CO8000 Cell Density, UV-Vis spectrophotometer D2800, biological sample digestion system, Nexus Gradient PCR thermal cycler, and Kjeldahl system KDN-04.

## 2.4. Method

### 2.4.1. Enrichment, isolation, and screening of a potential yeast strain

Five grams of each sample were homogenized in 45 mL of Yeast - Peptone - Dextrose broth (YPD broth). The suspension was incubated at 30°C with shaking at 200 rpm for 1 h to facilitate pre-activation. Subsequently, 10% (v/v) inoculum of the suspension was transferred into 45 mL of fresh YPD medium. This enrichment culture was maintained at 30°C, 200 rpm for 48 h to establish the 1<sup>st</sup> generation (T<sub>1</sub>). Subculturing was performed sequentially 3 times at 48 h intervals to obtain the 3<sup>rd</sup> generation (T<sub>3</sub>). The T<sub>3</sub>-enriched community was observed at 1000x magnification by crystal violet staining and subjected to isolation via the serial dilution method. 1 mL of the T<sub>3</sub> culture underwent ten-fold serial dilutions (up to 10<sup>-6</sup>) in 9 mL of 0.9% NaCl solution. Then, 100 µL of the culture was spread on YPD agar plates and incubated at 30°C for 48 h. Visible colonies were selected and purified by repeated streaking on YPD agar until pure cultures were obtained. The isolates were preserved in 30% glycerol at -20°C.


The suitability of yeast strains for SCP production is critically determined by their capacity for rapid biomass proliferation and high intracellular protein accumulation. Initially, 36 yeast strains were reactivated from -20°C storage on YPD agar plates at 30°C for 24 h. A loopful of activated cells was then inoculated into Erlenmeyer flasks containing 20 mL of YPD broth. Cultivation was performed at 30°C, 200 rpm for 24 h. To evaluate biomass production, the entire culture volume was centrifuged at 6000 rpm for 10 min. The cell pellet was collected and dried at 80°C until constant weight. The cell mass was used for total protein analysis by the Kjeldahl method.

### 2.4.2. DNA extraction and phylogenetic analysis

Genomic DNA was recovered through an integrated physicochemical cell lysis protocol [8]. Cell pellets were resuspended in lysis buffer fortified with 20%, followed by thermal incubation at 65°C for 120 min. Subsequently, 400 µL of Chloroform - Isoamyl alcohol was added, and the mixture was inverted thoroughly before centrifugation at 12,000 rpm for 10 min. The supernatant was collected and mixed with Isopropanol at a 1:1 ratio to precipitate the DNA, followed by incubation at -20°C for 1 h. The DNA pellet was recovered by centrifugation at 12,000 rpm for 10 min and subsequently resuspended in 50 µL of Milli-Q water. The ITS rDNA region was amplified via PCR using the primer pair ITS\_1 (5' TCCGTAGGTGAACCTGCGG 3') and ITS\_4 (5' TCCTCCGCTTATTGATATGC 3'), as described by White et al. (1989) [9]. The PCR program was detailed in **Figure 1**. The PCR samples were sequenced by Phusa Biochem Co., Ltd., and then were analyzed for homology against the NCBI GenBank database (<http://www.ncbi.nlm.nih.gov>) using the BLAST algorithm. A phylogenetic tree was constructed using MEGA 11 software based on the Neighbor-Joining method with 1000 bootstrap replications [10].

### 2.4.3. Effect of culture volume on YPI-Y2 biomass production

The effect of volume culture ratio was carried out with a range of 5 to 45 mL of YPD broth in a 100 mL conical flask. The seed culture of the YPI-Y2 strain was activated in YPD medium for 24h. After that, the seed culture was transferred to YPD broth with a different volume ratio to reach 1.0 value of OD<sub>600</sub>. Then the flasks were incubated at 30°C and shaken at 200 rpm. After 24 h, 5 mL of culture from each flask was centrifuged at 6000 rpm for 10 min. The resulting biomass was harvested and dried at 80°C until a constant weight was achieved. Biomass weight and cell density data were recorded. The experiments were performed in three independent replicates.



**Figure 1.** The detailed PCR program

#### 2.4.4. Optimization of nutritional components and culture conditions for growth and protein accumulation

Yeast cells YPI-Y2 were transferred into 45 mL YPD broth to establish the primary seed culture. Incubation was performed at 30°C with shaking at 200 rpm for 24 h. An appropriate volume of the seed culture was inoculated into 10 mL of the experimental media to achieve an  $OD_{600} = 1.0$ . This setup was designed to evaluate the effects on growth and concomitant protein accumulation across various parameters, including: 8 carbon sources (20 g/L of D-fructose, D-glucose, Xylose, Sucrose, Maltose, Maltose dextrin, Molasses and Starch soluble), 6 nitrogen sources (YP,  $(NH_4)_2SO_4$ ,  $(NH_4)_2HPO_4$ ,  $NH_4NO_3$ ,  $NH_4Cl$  and  $(NH_2)_2CO$  with total nitrogen of 3.8 g/L), 4 temperature points (25 - 40°C, increments 5°C), and 13 pH levels (3.0 - 9.0, increments 0.5). Key parameters, including biomass,  $OD_{600}$ , and total protein content, were quantified. The experiments were repeated three times.

#### 2.4.5. Analytical methods for biomass and total protein content

Cell dry weight (CDW) was determined according to the protocol described by Thuoc et al. (2014) [11]. Total protein content (TPC) was quantified using the Kjeldahl method [12, 13].


#### 2.4.6. Methodology for data analysis

Results were expressed as mean  $\pm$  standard deviation (SD) of 3 independent experiments, 164 conducted in triplicate. Data processing was performed using Microsoft Excel. Statistical significance was assessed using one-way ANOVA at a significance level of  $p < 0.05$ .


### 3. RESULTS AND DISCUSSION

#### 3.1. Enrichment, isolation, screening, and identification of yeast strains

The implementation of 3 consecutive enrichment cycles established an effective selective pressure, significantly suppressing non-adapted bacteria while fostering the dominance of yeast populations within the culture medium. The accumulation of ethanol, coupled with mild acidic conditions, acted as a natural competitive filter, thereby stabilizing the target microbial community. By the  $T_3$ , the microbial composition was notably refined, consisting predominantly of yeast species exhibiting robust growth kinetics and efficient substrate conversion capabilities (**Figure 3**). A total of 36 yeast strains were successfully isolated from five enriched communities.




**Figure 2.** Enrichment results of the five collected samples: (a) vegetable pickle, (b) “Tré” (fermented meat), (c) kimchi, (d) fermented grape, and (e) grape pomace




**Figure 3.** Microbial communities through successive enrichment processes of the vegetable pickle sample: (1) 1<sup>st</sup> generation, (2) 2<sup>nd</sup> generation, and (3) 3<sup>rd</sup> generation

After 24 h of cultivation, 15/36 strains (41.7%) yielded a CDW exceeding 10 g/L. The CDW values for this high-yield group ranged from 11 to 16 g/L. The highest CDW was observed in strains YFPS-M6 ( $15.6 \pm 0.42$  g/L) and YFPS-M10 ( $15.2 \pm 0.21$  g/L) (**Figure 4**). Of these, 3 strains were isolated from Kimchi and exhibited CDW values of approximately 11 g/L. Among the remaining isolates, 15 strains displayed CDW values between 6 and 10 g/L, while 8 strains produced less than 6 g/L. In a study by Trang et al. (2017), the *S. cerevisiae* SC2.75 strain achieved a CDW of 10.71 g/L after 18h of cultivation [4]. These comparisons indicate that the 15 effectively regulate metabolism to maximize biomass production, even under non-optimized conditions.



**Figure 4.** CDW and TPC values of promised yeast strains



**Figure 5.** Phylogenetic tree of yeast strains based on ITS rDNA sequences BLASTed on NCBI. A - YPI-Y2 relationship with the *Pichia* genus; B - Other strain relationship with the *Candida* genus


TPC analysis of the ten yeast strains with the highest biomass revealed significant variation in SCP production capacity, with values ranging from 57 to 62.3% (g/100 g of CDW). Strains named YFPS-M5, YFPS-M6, YFPS-M7, YFPS-M8, and YFPS-M10 could produce relatively high cell mass ranging from 13 to 15 g/L, but TPC values achieved were not superior. Conversely, strains YPI-Y2 and YKCS-Y3 exhibited outstanding protein accumulation capacity with TPC value at 62.3% and 61.5%, respectively (**Figure 4**). A study by Rachamontree et al. (2015) reported that the strain *P. kudriavzevii* MSY-2 achieved a TPC of 66.8% (g/100 g of biomass) only under optimized culture conditions [14]. Remarkably, several strains in this study achieved TPC values comparable to this benchmark without any prior optimization. This finding underscores the robust physiological potential of these isolates for efficient SCP production.

Predominant yeast strains were successfully identified to the genus level based on ITS sequence analysis. The constructed phylogenetic tree showed that the yeast strain designated YPI-Y2 belongs to the genus *Pichia* with a similarity level of 99.77% to the species *P. kudriavzevii* (**Figure 5A**). The other strains were all identified as belonging to the genus *Candida* with a similarity level of over 99% to the species *C. tropicalis* (**Figure 5B**). Strain YPI-Y2 was selected as the best candidate for further investigation.

### 3.2. Culture characteristics of the YPI-Y2 strain for biomass production and protein accumulation

#### 3.2.1. Effect of culture volume ratio


The investigation across 8 different culture volumes revealed that the V5 and V10 conditions supported the most favorable growth. Specifically, the maximum optical OD<sub>600</sub> of 19.7 was recorded at V10. Correspondingly, the CDW at V10 (13.7 g/L) surpassed that of V5 (12.4 g/L). Notably, a progressive decline in growth was observed as the volume increased from V15 to V40. The lowest performance was recorded at V40, with an OD<sub>600</sub> of 12.95 and a CDW of only 7.2 g/L (**Figure 6**). These findings demonstrate a critical correlation between the medium volume and the liquid-air interface area, which directly influences the O<sub>2</sub> transfer rate and consequently, the biomass productivity of the yeast strain. This observation aligns with the study by Nayana et al. (2023), which corroborated that culture volume and surface area availability significantly impact the yield and physiological activities of aerobic microorganisms due to O<sub>2</sub> limitation [15]. Based on these results, a medium to flask volume ratio of 1/10 was selected as the ideal condition for further characterization of the strain.



**Figure 6.** Effect of culture volume ratio on CDW and OD<sub>600</sub> values of yeast strain YPI-Y2


#### 3.2.2. Effect of pH on growth and protein accumulation

The TPC of the yeast strain exhibited significant fluctuations in response to variations in the initial pH of the culture medium. As the pH increased from 3.0 to 4.0, the TPC remained relatively stable within the range of 51% to 52% (**Figure 7**). This observation indicates that the strain retains its capacity for protein biosynthesis even under mildly acidic conditions. The obtained results, within the pH range of 5.0 to 6.5, a progressive increase in protein content was observed, reaching a peak at pH 6.5 (64.75%). This peak coincided with the maximal values recorded for OD<sub>600</sub> of 19.65 and CDW of 13.63 g/L. Under relatively acidic pH conditions (pH 3.0), the CDW value of strain YPI-Y2 reached 88.6% compared to the optimal pH point (pH 6.5). The results are quite similar to those of Thu et al. (2023) and Xu et al. (2024) for *P. kudriavzevii* [16, 17].



**Figure 7.** Effect of initial pH on growth and protein accumulation

### 3.2.3. Effect of temperature




**Figure 8.** Effect of culture temperature on growth and protein accumulation

The results indicate that incubation temperature exerts a significant influence on both the growth capacity and protein accumulation of the yeast strain. A substantial increase in protein content was observed as the temperature rose from 25 to 30°C. The value peaked at 65.44% at 30°C, coinciding with maximal OD<sub>600</sub> of 19.95 and CDW of 13.45 (Figure 8). This temperature was identified as the optimum, enabling cells to maintain a high metabolic rate and robust protein synthesis.

Publications on the yeast *P. kudriavzevii* show similar results to the impact of temperature on growth as in this study. Nieto-Sarabia et al. (2022) showed a wide temperature range for the growth of strain *P. kudriavzevii* 4A and best exhibition at 30 - 40°C [18]. In the study of Chamnipa et al. (2018), the strain *P. kudriavzevii* RZ8-1 could grow well in the temperature range of 30 - 40°C and slightly decreased at 42°C, but sharply at 45°C [19].

### 3.2.4. Effect of carbon sources

Carbon serves as a fundamental macronutrient for all living organisms. Furthermore, it functions as the primary energy source driving microbial growth and the concomitant accumulation of associated secondary products. In this study, various carbon sources were investigated at a concentration of 2% (w/v) with the absence of a carbon source as a negative control to evaluate the metabolic performance of yeast strain YPI-Y2 using minimal salt medium (MSM) [20]. The results indicated that strain YPI-Y2 was capable of assimilating 7/8 tested carbon sources for growth, while maintaining a state of survival on the remaining source (xylose).




**Figure 9.** Effect of carbon sources on growth and protein accumulation of yeast strain YPI-Y2

Among the assimilable sources, YPI-Y2 exhibited superior growth and protein accumulation when cultivated on D-fructose and D-glucose. Specifically, maximal values for CDW, OD600, and TPC were recorded with D-fructose (13.3 g/L, 19.05, and 65.35%, respectively). In comparison, D-glucose yielded slightly lower values of 12.8 g/L, 18.8, and 59.39% (**Figure 9**). These findings suggest that D-fructose is metabolized more efficiently than D-glucose by this specific strain. Compared to monosaccharides, complex carbon sources (sucrose, maltose, maltodextrin, and starch) resulted in significantly reduced metabolic efficiency. This reduction is attributed to their complex chemical structures (polymers, oligosaccharides), which necessitate additional energy expenditure for hydrolysis before assimilation. This energy diversion limits the resources available for cellular proliferation and protein synthesis. Consequently, the average OD600 for this group ranged from 8.82 to 10.08, approximately 1.4-fold higher than the control but 2.03-fold lower than that of D-fructose. Similarly, CDW values ranged from 3.2 to 4.7 g/L (4.1-fold lower than D-fructose), and TPC values were moderate, ranging from 54.9% to 58.3%. Molasses, a complex substrate containing various sugars, minerals, and potential inhibitors that may induce cellular stress or inhibit hydrolytic enzymes, supported moderate growth. Strain YPI-Y2 demonstrated a metabolic capability on molasses second only to the monosaccharides, achieving an OD600 of 15.17, CDW of 8.2 g/L, and TPC of 58.34%. High total protein content and biomass due to molasses utilization is an advantage of the YPI-Y2 strain by solving an inexpensive by-product and thus increasing the competitiveness at an industrial production scale. Despite its ability to utilize complex sugars, YPI-Y2 failed to assimilate xylose effectively. This limitation is likely due to the absence of key enzymes, such as xylose reductase and xylitol dehydrogenase, which are required to convert xylose into xylulose for entry into the Pentose Phosphate Pathway [21, 22]. Our results align with those of Nava et al. (2017), demonstrating that *P. kudriavzevii* ITV-S42 preferentially metabolized glucose and fructose as primary carbon sources, with negligible xylose catabolism and limited sucrose utilization due to deficient metabolic enzymes [23]. Notably, Koutinas et al. (2016) reported that *P. kudriavzevii* KVMP10 possessed efficient xylose-fermenting capacity at elevated temperatures [24], indicating genetic diversity within this species. This finding highlights genetic diversity among different *P. kudriavzevii* strains. The OD600 of 6.23 and CDW of 3.05 g/L on xylose showed minimal deviation from the negative control (OD600 of 6.68; CDW of 2.48 g/L). However, the TPC remained relatively high at 59.96%. This suggests that while xylose does not support proliferation, a fraction may be partially metabolized to provide maintenance energy for organelle function and cellular survival.

### 3.2.5. Effect of nitrogen sources

Nitrogen is an indispensable element for biological functions, serving as a fundamental constituent of vital cellular macromolecules, including protein, nucleic acids, and ATP... Consequently, it plays a critical governing role in microbial growth. To evaluate the growth and concomitant protein accumulation of yeast strain YPI-Y2, six distinct nitrogen sources were investigated: combined Yeast-Peptone (YP),  $(\text{NH}_4)_2\text{SO}_4$ ,  $(\text{NH}_4)_2\text{HPO}_4$ ,  $\text{NH}_4\text{NO}_3$ ,  $\text{NH}_4\text{Cl}$ , and  $(\text{NH}_2)_2\text{CO}$ , with the absence of nitrogen source as a negative control.



**Figure 10.** Effect of nitrogen sources on growth and protein accumulation of yeast strain YPI-Y2

The results indicated that strain YPI-Y2 exhibited optimal performance when cultivated on a yeast-peptone combination (YP), the organic nitrogen source. Specifically, the TPC reached 66.62%, while the maximal OD<sub>600</sub> and CDW peaked at 19.08 and 16.15 g/L, respectively (**Figure 10**). Compared to other groups, cultivation on YP resulted in a 4-folds increase in cell density (OD<sub>600</sub>) and a 2.6-folds increase in protein accumulation. This superiority is likely attributable to the presence of readily absorbable amino acids and peptides in peptone and yeast extract. These components not only provide a rapid nitrogen supply but also act as precursors that stimulate the synthesis of intracellular structural proteins and metabolic enzymes. Xu et al. (2022) reported that the *P. kudriavzevii* XTY1 strain demonstrated efficient utilization of organic nitrogen during heterotrophic nitrification, which is consistent with our findings showing that organic nitrogen utilization capacity by *P. kudriavzevii* YPI-Y2 exhibited superior performance relative to inorganic nitrogen sources [21]. Among the inorganic sources, (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub> (53.44%), (NH<sub>4</sub>)<sub>2</sub>HPO<sub>4</sub> (51.56%), and NH<sub>4</sub>Cl (52.11%) yielded superior efficiency compared to ammonium nitrate and urea. These salts provide readily metabolizable ammonium ions, NH<sub>4</sub><sup>+</sup>, and facilitate pH stability, thereby supporting moderate biomass production (CDW ranging from 5.7 to 7.6 g/L) and enhancing protein accumulation. NH<sub>4</sub>NO<sub>3</sub> in a lower TPC of 48.36%. This reduction is likely because the nitrate ion NO<sub>3</sub><sup>-</sup> requires energy-intensive reduction before assimilation, effectively slowing down the rate of nitrogen incorporation. Although urea supported relatively high OD<sub>600</sub> and CDW values, the total protein content reached only 35.62%. This disparity suggests that the rate of urea hydrolysis into utilizable NH<sub>4</sub><sup>+</sup> is slower compared to the direct uptake available from ammonium salts. Due to low cost and availability, ammonium could be potentially important nitrogen when using the YPI-Y2 strain in single cell protein production at an industrial scale.

#### 4. CONCLUSION

The potential yeast strains (36 strains) were isolated from fermented food that exhibited good growth and high protein accumulation, ranging from 57% to 62.3% (g/100 g cell biomass). Among them, the most promising strain, *P. kudriavzevii* YPI-Y2, was selected for further investigation. The optimal conditions for the growth and concomitant protein accumulation of YPI-Y2 were determined as follows: an ideal culture volume ratio of 10%, D-fructose (20 g/L), and a combination of yeast-peptone (YP, 3.8 g/L) as the carbon and nitrogen sources, pH of 6.5, and temperature of 30°C. Under these optimal conditions, the average CDW and TPC reached 16.12 g/L and 66.62% (g/100 g biomass), respectively, within 24 h of cultivation. These results substantiate YPI-Y2's superior protein accumulation capacity and short fermentation duration, demonstrating its promising potential for industrial single-cell protein production. Further studies on safety and fermentation technology should be conducted to provide robust data for the application of the YPI-Y2 strain for single cell protein production at an industrial scale.

#### ACKNOWLEDGEMENT

This study was conducted with the support of Bioresource Research Center, Phenikaa School of Engineering, Phenikaa University.

#### REFERENCES

- [1]. Y. P. Li, F. Ahmadi, K. Kariman *et al.*, "Recent advances and challenges in single cell protein (SCP) technologies for food and feed production," *npj Science of Food*, vol. 8, 2024.
- [2]. M. E. Jach, A. Serefko, M. Ziaja, and M. Kieliszek, "Yeast protein as an easily accessible food source," *Metabolites*, vol. 12, no. 1, 2022.
- [3]. F. Samiksha, E. M. D. San Valentin, G. Li, M. Blazer, T. S. McCormick, and M. Ghannoum, "A narrative review on the functional applications, safety, and probiotic characteristics of *Pichia*," *Nutrients*, vol. 17, 2025.
- [4]. T. H. T. Ngo and V. H. Vu, "Study on optimal fermentation conditions for the high biomass production of *Saccharomyces cerevisiae* SC2.75," *Vietnam Journal of Biotechnology*, vol. 15, no. 3, pp. 581-588, 2017 (in Vietnamese).
- [5]. T. M. H. Pham, T. T. H. Nguyen, and N. U. Le, "Production of *Saccharomyces cerevisiae* biomass from *Ulva lactuca* algae hydrolysis and experiments for use in commercial Pacific oyster farming," *Journal of Science of Can Tho University*, vol. 60, Special issue SDMD, pp. 138-145, 2024 (in Vietnamese).

- [6]. Ministry of Industry and Trade, "Utilizing by-products of the paper industry to produce animal feed," 2025. [Online]. Available: <https://nq57.igip.gov.vn/tin-tuc/t3207>. [Accessed: Dec. 26, 2025] (in Vietnamese).
- [7]. Y. Chu *et al.*, "Advances in the application of the non-conventional yeast *Pichia kudriavzevii* in food and biotechnology industries," *Journal of Fungi*, vol. 9, no. 170, 2023.
- [8]. L. E. Epperson and M. Strong, "A scalable, efficient, and safe method to prepare high quality DNA from mycobacteria and other challenging cells," *Journal of Clinical Tuberculosis and Other Mycobacterial Diseases*, vol. 19, 2020.
- [9]. T. J. White, T. D. Bruns, S. B. Lee, and J. W. Taylor, "Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics," in *PCR Protocols: A Guide to Methods and Applications*, M. A. Innis, D. H. Gelfand, J. J. Sninsky, and T. J. White, Eds. New York, NY, USA: Academic Press, 1990, pp. 315–322.
- [10]. L. Newman, A. L. J. Duffus, and C. Lee, "Using the free program MEGA to build phylogenetic trees from molecular data," *American Biology Teacher*, vol. 78, no. 7, pp. 608-612, 2016.
- [11]. D. Van-Thuoc and J. Quillaguamán, "Improving culture conditions for poly(3-hydroxybutyrate-co-3-hydroxyvalerate) production by *Bacillus* sp. ND153, a bacterium isolated from a mangrove forest in Vietnam," *Annals of Microbiology*, vol. 64, pp. 991-997, 2014.
- [12]. Vietnam Institute of Quality Standards, *Food and animal feed - General guidance on nitrogen content determination by Kjeldahl method*, TCVN 10034:2013 (ISO 1871:2009), 2013 (in Vietnamese).
- [13]. AOAC International, "AOAC Official Method 2001.11: Protein (Crude) in Animal Feed, Forage, Grain, and Oilseeds - Block Digestion Method Using Copper Catalyst and Steam Distillation into Boric Acid," in *Official Methods of Analysis of AOAC International*, 2023.
- [14]. P. Rachamontree, T. Phusantisampan, N. Woravutthikul, P. Pornwongthong, and M. Sriariyanun, "Selection of *Pichia kudriavzevii* strain for the production of single-cell protein from cassava processing waste," *International Journal of Biological, Food, Veterinary and Agricultural Engineering*, vol. 9, no. 5, 2015.
- [15]. K. Nayana, D. Vidya, K. Soorya *et al.*, "Effect of volume and surface area on growth and productivity of microalgae in culture system," *BioEnergy Research*, vol. 16, pp. 1013-1025, 2022.
- [16]. Q. Xu, W. Huang, Y. Li *et al.*, "Isolation and identification of epiphytic *Pichia kudriavzevii* from loquat peels and investigation of its fermentation characteristics for liquor production," *Archives of Microbiology*, vol. 206, no. 1, 2024.
- [17]. N. T. T. Thu *et al.*, "Evaluation of polyhydroxyalkanoate (PHA) synthesis by *Pichia* sp. TSL24 yeast isolated in Vietnam," *Scientific Reports*, vol. 13, 2023.
- [18]. V. L. Nieto-Sarabia, C. B. Ballinas-Cesatti, G. Melgar-Lalanne, E. Cristiani-Urbina, and L. Morales-Barrera, "Isolation, identification, and kinetic and thermodynamic characterization of a *Pichia kudriavzevii* yeast strain capable of fermentation," *Food and Bioprocess Processing*, vol. 131, pp. 109-124, 2022.
- [19]. N. Chamnipa, S. Thanonkeo, P. Klanrit, and P. Thanonkeo, "The potential of the newly isolated thermotolerant yeast *Pichia kudriavzevii* RZ8-1 for high-temperature ethanol production," *Brazilian Journal of Microbiology*, vol. 49, no. 2, pp. 378-391, 2018.
- [20]. T. Xu *et al.*, "Fungus *Pichia kudriavzevii* XTY1 and heterotrophic nitrifying bacterium *Enterobacter asburiae* GS2 cannot efficiently transform organic nitrogen via hydroxylamine and nitrite," *Frontiers in Microbiology*, vol. 13, 2022.
- [21]. M. Jeppsson, B. Johansson, B. Hahn-Hägerdal, and M. F. Gorwa-Grauslund, "Reduced oxidative pentose phosphate pathway flux in recombinant xylose-utilizing *Saccharomyces cerevisiae* strains improves the ethanol yield from xylose," *Applied and Environmental Microbiology*, vol. 68, no. 4, pp. 1604-1609, 2002.
- [22]. Z. Jiang *et al.*, "Glucose-6-phosphate dehydrogenases: The hidden players of plant physiology," *International Journal of Molecular Sciences*, vol. 23, no. 24, 2022.
- [23]. L. E. Díaz-Nava, N. Montes-García, J. M. Domínguez, and M. G. Aguilar-Uscanga, "Effect of carbon sources on the growth and ethanol production of native yeast *Pichia kudriavzevii* ITV-S42 isolated from sweet sorghum juice," *Bioprocess and Biosystems Engineering*, vol. 40, no. 7, pp. 1069-1077, 2017.
- [24]. M. Koutinas, M. Patsalou, S. Stavrinou, and I. Vyrides, "High temperature alcoholic fermentation of orange peel by the newly isolated thermotolerant *Pichia kudriavzevii* KVMP10," *Letters in Applied Microbiology*, vol. 62, no. 1, pp. 75-83, 2016.